

Figure 1: Various synthetic pathways for the biosynthesis of DHA (docosahexaenoic acid)




Figure 2: Substrate specificity of the  $\Delta 5$ -elongase (SEQ ID NO: 53) with regard to different fatty acids





Figure 3: Reconstitution of DHA biosynthesis in yeast starting from 20:5 $\omega$ 3.



Figure 4: Reconstitution of DHA biosynthesis in yeast starting from 18:4 $\omega$ 3.

Figure 5: Fatty acid composition (in mol%) of transgenic yeasts which had been transformed with the vectors pYes3-OmELO3/pYes2-EgD4 or pYes3-OmELO3/pYes2-EgD4+pESCLeu-PtD5. The yeast cells were cultured in minimal medium without tryptophan and uracil/ and leucin in the presence of 250  $\mu$ M 20:5 $^{\Delta 5,8,11,14,17}$  and 18:4 $^{\Delta 6,9,12,15}$ , respectively. The fatty acid methyl esters were obtained from cell sediments by acid methanolysis and analyzed via GLC. Each value represents the mean (n=4)  $\pm$  standard deviation.

| Fatty acids                      | pYes3-OmELO/pYes2-EgD4                   | pYes3-OmELO/pYes2-EgD4 EgD4<br>+ pESCLeu-PtD5 |
|----------------------------------|--|---|
|                                  | Feeding of 20:5 $^{\Delta 5,8,11,14,17}$ | Feeding of 18:4 $^{\Delta 6,9,12,15}$         |
| 16:0                             | 9.35 $\pm$ 1.61                          | 7.35 $\pm$ 1.37                               |
| 16:1 $^{\Delta 9}$               | 14.70 $\pm$ 2.72                         | 10.02 $\pm$ 1.81                              |
| 18:0                             | 5.11 $\pm$ 1.09                          | 4.27 $\pm$ 1.21                               |
| 18:1 $^{\Delta 9}$               | 19.49 $\pm$ 3.01                         | 10.81 $\pm$ 1.95                              |
| 18:1 $^{\Delta 11}$              | 18.93 $\pm$ 2.71                         | 11.61 $\pm$ 1.48                              |
| 18:4 $^{\Delta 6,9,12,15}$       | -  | 7.79 $\pm$ 1.29                               |
| 20:1 $^{\Delta 11}$              | 3.24 $\pm$ 0.41                          | 1.56 $\pm$ 0.23                               |
| 20:1 $^{\Delta 13}$              | 11.13 $\pm$ 2.07                         | 4.40 $\pm$ 0.78                               |
| 20:4 $^{\Delta 8,11,14,17}$      | -  | 30.05 $\pm$ 3.16                              |
| 20:5 $^{\Delta 5,8,11,14,17}$    | 6.91 $\pm$ 1.10                          | 3.72 $\pm$ 0.59                               |
| 22:4 $^{\Delta 10,13,16,17}$     | -  | 5.71 $\pm$ 1.30                               |
| 22:5 $^{\Delta 7,10,13,16,19}$   | 8.77 $\pm$ 1.32                          | 1.10 $\pm$ 0.27                               |
| 22:6 $^{\Delta 4,7,10,13,16,19}$ | 2.73 $\pm$ 0.39                          | 0.58 $\pm$ 0.10                               |

Figure 6: Feeding experiment for determining the functionality and substrate specificity with yeast strains






Figure 7: Elongation of eicosapentaenoic acid by Otlol1






Figure 9: Expression of TpELO1 in yeast




Figure 10: Expression of TpELO3 in yeast




Figure 11: Expression of *Thraustochytrium*  $\Delta 5$ -elongase TL16/pYES2.1 in yeast




Figure 12: Desaturation of  $\gamma$ -linolenic acid (18:2  $\omega$ 6-fatty acid) to give  $\alpha$ -linolenic acid (18:3  $\omega$ 3-fatty acid) by Pi-omega3Des.




Figure 13: Desaturation of  $\gamma$ -linolenic acid (18:2  $\omega$ 6-fatty acid) to give stearidonic acid (18:4  $\omega$ 3-fatty acid) by Pi-omega3Des.




Figure 14: Desaturation of C20:2  $\omega$ 6-fatty acid to give C20:3  $\omega$ 3-fatty acid by Pi- $\omega$ 3Des.




Figure 15: Desaturation of C20:3  $\omega$ 6-fatty acid to give C20:4  $\omega$ 3-fatty acid by Pi- $\omega$ 3Des.




Figure 16: Desaturation of arachidonic acid (C20:4  $\omega$ 6-fatty acid) to give eicosapentaenoic acid (C20:5  $\omega$ 3-fatty acid) by Pi-omega3Des.






Figure 17: Desaturation of docosatetraenoic acid (C22:4  $\omega$ 6-fatty acid) to give docosapentaenoic acid (C22:5  $\omega$ 3-fatty acid) by Pi-omega3Des.




Figure 18: Substrate specificity of Pi-omega3Des with regard to different fatty acids




Figure 19: Desaturation of phospholipid-bound arachidonic acid to give EPA by Pi-Omega3Des





Figure 20: Conversion of linoleic acid (arrow) to give  $\gamma$ -linolenic acid ( $\gamma$ -18:3) by Ot-Des6.1.

Absorption mAU



Retention time

Figure 21: Conversion of linoleic acid and  $\alpha$ -linolenic acid (A and C), and reconstitution of the ARA and EPA synthetic pathways, respectively, in yeast (B and D) in the presence of OtD6.1.




Figure 22: Expression of ELO(XI) in yeast

Absorption in mA






Figure 23:




Figure 24: Elongation of eicosapentaenoic acid by OtElo1 (B) and OtElo1.2 (D), respectively. The controls (A, C) do not show the elongation product (22:5 $\omega$ 3).






Figure 25: Elongation of arachidonic acid by OtElo1 (B) and OtElo1.2 (D), respectively. The controls (A, C) do not show the elongation product (22:4 $\omega$ 6).




Figure 26: Elongation of 20:5n-3 by the elongases At3g06470.




Figure 27: Substrate specificity of the *Xenopus* Elongase (A), *Ciona* Elongase (B) and *Oncorhynchus* Elongase (C)




Figure 28: Substrate specificity of the *Ostreococcus*  $\Delta 5$ -elongase (A), the *Ostreococcus*  $\Delta 6$ -elongase (B), the *Thalassiosira*  $\Delta 5$ -elongase (C) and the *Thalassiosira* *Ostreococcus*  $\Delta 6$ -elongase (D)





Figure 29: Expression of the *Phaeodactylum tricornutum*  $\Delta 6$ -elongase (PtELO6) in yeast. A) shows the elongation of the C18:3 $^{\Delta 6,9,12}$  fatty acid and B) the elongation of the C18:3 $^{\Delta 6,9,12,15}$  fatty acid

A)



B)




Figure 30: Figure 30 shows the substrate specificity of PtELO6 with regard to the substrates fed.




Figure 31: Gas-chromatographic analysis of the seed of a transgenic plant, transformed with pSUN-5G.




Figure 32: Gas-chromatographic analysis of the seed of a transgenic plant, transformed with pGPTV-D6Des(Pir)\_D5Des(Tc)\_D6Elo(PP)\_12Des(Co)






Figure 33: DHA in transgenic seeds of *Brassica juncea*. The plants were transformed with the construct pSUN-8G.

